

Pro LEBANON Scape®

FERTILIZER with LockUp™ Herbicide

19-0-6



MESA

GUARANTEED ANALYSIS

Total Nitrogen (N)..... 19%
 2.6% Ammoniacal Nitrogen
 1.2% Water Insoluble Nitrogen*
 11.8% Urea Nitrogen
 2.8% Other Water Soluble Nitrogen*
 Soluble Potash (K₂O)..... 6%
 Sulfur (S)..... 3.0%
 3.0% Combined Sulfur (S)



F659 **22-54361**

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash,
 Chlorine (Cl) not more than 6.0%
 *4.6% Slowly available Nitrogen from Methylene Ureas.

SUGGESTED SPREADER SETTINGS

Spreader	19-0-6		19-0-6	
	100lb/acre 2,461,000 sq. ft.	175lb/acre 4,011,000 sq. ft.	100lb/acre 2,461,000 sq. ft.	175lb/acre 4,011,000 sq. ft.
LebanonTurf	4"	4"	Leuca (Lebanon Herbicide) 1/17	1/21
Lebanon AcePro	2"	4"	ProScape 19-0-6	M
Earthway Rotary	14	10	ProScape 19-0-6	M
Casey	25	20	Sprayer	40
Lab	5.0	0.0	Vicon (Lebanon Herbicide)	24

Note: These settings are provided as a suggested starting point in calculating each fertilizer application. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by seller or the spreader manufacturer.

For postemergence control of annual and perennial broadleaf weeds in established turfgrass, including residential lawns, golf courses, parks, sports fields, sod farms and around commercial buildings and other recreational and commercial turfgrass areas

ACTIVE INGREDIENT:
 2,4-dichlorophenoxyacetic acid..... 1.04%
 penoxsodium 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy[1,2,4]triazolo[1,5-c]pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide... 0.01%
 dicamba 3,6-dichloro-o-anisic acid..... 0.08%
OTHER INGREDIENTS:..... 98.87%
Total..... 100.00%

Acid equivalent: 2,4-D acid - 0.86% - 17.2 lb/ton
 dicamba - 0.08% - 1.5 lb/ton

EPA Reg. No. 22710-569-001 EPA Est. 561-PA-01, 95141-01
 ProScape and MESA are registered trademarks of Lebanon Seaboard Corporation. Supercept used is first letter of last number.

KEEP OUT OF REACH OF CHILDREN CAUTION

Precautionary Statements Hazards to Humans and Domestic Animals

Harmful if Absorbed Through Skin, If Inhaled, Or If Swallowed • Causes Moderate Eye Irritation

Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wear protective eyewear, long-sleeved shirt, long pants, socks, shoes, and chemical-resistant gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

	First Aid
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
If swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may contact 1-800-969-5954 for emergency medical treatment information.

Doc Id: 526509

ACCEPTED
VIA NOTIFICATION
LABEL NOT REVIEWED

May 4, 2011

New York State Department
of Environmental Conservation
Division of Materials Management
Pesticide Product Registration



Distributed by:
Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042
Made in the U.S.A.

NET WEIGHT **40** LBS. (18.14 Kg)

FERTILIZER

with LockUp™ Herbicide

19-0-6



MESA

P A T E N T E D

GUARANTEED ANALYSIS

Total Nitrogen (N) 19%
 2.6% Ammoniacal Nitrogen
 1.8% Water Insoluble Nitrogen*
 11.8% Urea Nitrogen
 2.8% Other Water Soluble Nitrogen*

Soluble Potash (K₂O) 6%
Sulfur (S) 3.0%
 3.0% Combined Sulfur (S)



Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.
 Chlorine (Cl) not more than 6.0%
 *4.6% Slowly available Nitrogen from Methylene Ureas.

SUGGESTED SPREADER SETTINGS

Spreader	150#/acre		175#/acre		Spreader	150#/acre		175#/acre	
	3.4#/1,000 sq. ft.		4#/1,000 sq. ft.			3.4#/1,000 sq. ft.		4#/1,000 sq. ft.	
LebanonTurf	4½	4¾	Lesco (letter/Numeric)	1/17	J/21				
Andersons AccuPro	K	M	PennMulch HVO	K	M				
Earthway Rotary	14	16	ProScape SS	K	M				
Gandy	26	28	Spyker	4½	4¾				
Lely	5 II	6 II	Vicon (all models)	24	28				

Note: These settings are provided as a suggested starting point in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by seller or the spreader manufacturer.

For postemergence control of annual and perennial broadleaf weeds in established turfgrass, including residential lawns, golf courses, parks, sports fields, sod farms and around commercial buildings and other recreational and commercial turfgrass areas

ACTIVE INGREDIENT:
 2,4-dichlorophenoxyacetic acid..... 1.04%
 penoxsulam: 2-(2,2-difluoroethoxy)-N-(5,8-dimethoxy(1,2,4-triazolo(1,5-c)pyrimidin-2-yl)-6-(trifluoromethyl)benzenesulfonamide.. 0.01%
 dicamba: 3,6-dichloro-o-anisic acid..... 0.08%
OTHER INGREDIENTS:..... 98.87%
Total 100.0%

Acid equivalent: 2,4-D acid - 0.86% - 17.2 lb/ton
 dicamba - 0.08% - 1.5 lb/ton

EPA Reg. No. 62719-590-961
 ProScape and MESA are registered trademarks of Lebanon Seaboard Corporation.
 EPA Est. 961-PA-01; 961-IL-01
 Superscript used is first letter of lot number.

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

**Precautionary Statements
 Hazards to Humans and Domestic Animals**

Harmful If Absorbed Through Skin, If Inhaled, Or If Swallowed • Causes Moderate Eye Irritation

Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wear protective eyewear, long-sleeved shirt, long pants, socks, shoes, and chemical-resistant gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

First Aid


If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may contact 1-800-992-5994 for emergency medical treatment information.



Distributed by:
Lebanon Seaboard Corporation
 1600 E. Cumberland St. • Lebanon, PA 17042
 Made in the U.S.A.

NET WEIGHT 40 LBS. (18.14 KG)

Personal Protective Equipment (PPE)

All mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems and enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)].

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark except as noted on appropriate labels. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

2,4-D has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted entry interval, and notification to workers (as applicable). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls worn over short-sleeve shirt and short pants
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear for overhead exposure
- Protective eyewear

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated area.

Non-Agricultural Use Requirements

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: Do not enter or allow people (or pets) to enter area until dusts have settled.

Storage and Disposal

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool dry place in original container.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on the label. If terms are unacceptable, return at once unopened.

General Information

ProScape fertilizer with LockUp™ herbicide is a stable combination of three herbicides. It provides postemergence control of annual, biennial and perennial broadleaf weeds in established turfgrass, including residential lawns, golf courses, parks, sports fields, sod farms and around commercial buildings and other recreational and commercial turfgrass areas.

General Use Precautions and Restrictions

- Do not apply ProScape fertilizer with LockUp™ on dichondra, carpetgrass, St. Augustinegrass, bahiagrass, or to golf course putting greens or tees, or turfgrass containing desirable clovers.
- Do not apply ProScape fertilizer with LockUp™ directly to, or otherwise permit ProScape fertilizer with LockUp™ to come into contact with, flowers, ornamental shrubs or trees, or other desirable broadleaf plants, as serious plant injury may occur.
- Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at application.
- The active ingredient in this product may leave residues in treated grass clippings which could be harmful to some plants and trees. Do not collect treated grass clippings for use as mulch around plants. The applicator must notify in writing the appropriate residences/individuals of this precaution when this product is applied by a lawn service/commercial applicator.
- To minimize the potential for turfgrass injury, allow four weeks between applications.
- Do not water, mow or disturb treated area for 48 hours after application.
- Use ProScape fertilizer with LockUp™ on solid (monoculture) stands of perennial ryegrass or tall fescue at a maximum rate of 175 lb per acre. Some yellowing of these turfgrasses may occur within two weeks of application. Any discoloration is transitory and will dissipate under adequate growing conditions.
- Apply ProScape fertilizer with LockUp™ to newly seeded or sprigged turfgrass only after the third mowing or after the turfgrass seedlings have developed an adequate secondary root system. Reseeding is not recommended for at least three to four weeks after application.
- Do not apply more than 175 lb of ProScape fertilizer with LockUp™ per acre per year.
- Do not make more than two applications of ProScape fertilizer with LockUp™ per acre per year.
- The re-treatment interval is 21 days for applications to turfgrass grown for sod or seed.
- Florida Applications: For application to sports turf (non-agricultural land planted exclusively for golf courses, parks and athletic fields), follow "SL191, "Recommendations for N, P, K and Mg for Golf Course & Athletic Field Fertilization Base on Mehlich I Extractant" available at <http://edis.ifas.ufl.edu/SS404>, or "BMP's for Enhancement of Environmental Quality on Florida's Golf Courses" available at <http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/qtlbmp07.pdf>.

Treatment on Turfgrass Species Not Listed on the Label for ProScape fertilizer with LockUp™

Users who wish to use ProScape fertilizer with LockUp™ on turfgrass species not listed on this label may determine the suitability for use by treating a small area at a specified rate. Prior to treatment of larger areas, the treated area should be observed for any sign of herbicidal injury during 30 days of normal growing conditions to determine if the target turfgrass species is tolerant. Applications should not be made to turfgrass under stress. The user assumes responsibility for any plant damage or other liability resulting from use of ProScape fertilizer with LockUp™ on turfgrass species not listed on this label.

ProScape fertilizer with LockUp™ is recommended for use on the following established turfgrass species (numbers in parentheses (-) refer to footnotes below):

Established Cool Season Turfgrass Common Name	Scientific Name	Established Warm Season Turfgrass Common Name	Scientific Name
annual bluegrass	Poa annua	bermudagrass (common or hybrid)	Cynodon dactylon
bluegrass, Kentucky	Poa pratensis	centipedegrass (2)	Eremochloa ophiuroides
fescue, chewing	Festuca rubra var. commutata	fescue, tall (growing in warm season areas) (1)	Festuca arundinaceae
fescue, creeping red	Festuca rubra	kikuyugrass	Pennisetum clandestinum
fescue, sheeps	Festuca ovina	zoysiagrass	Zoysia japonica
fescue, tall (1)	Festuca arundinaceae	zoysiagrass	Zoysia tenuifolia
ryegrass, perennial (1)	Lolium perenne		

1. Use ProScape fertilizer with LockUp™ on these species at a maximum rate of 175 lb (0.02 lb ai peroxsulum) per acre per application and only if some injury can be tolerated. A slight yellowing of turfgrass may occur within two weeks of application. Any discoloration is transient and is generally gone within four weeks of treatment under adequate growing conditions.
2. Do not apply to centipedegrass during the period of winter dormancy. Grass should be green and actively growing at application.

Weeds Controlled or Suppressed

Generally, application rates at the lower end of the rate range will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult (plant stress conditions such as drought or extreme temperatures, dense weed stands and/or larger weeds), the higher rate within the rate range or repeat applications will be needed. Weeds growing in the absence of competition from other vegetation generally require higher rates or repeat applications to obtain satisfactory control or suppression.

Weeds Controlled or Suppressed

bedstraw, burdock, chickweed, chicory, clover, dandelion, dock, dollarweed, eclipta, healall, ivy (ground), knotweed, lambsquarters, medic (black), morningglory, peppergrass, pigweed, plantain, plantain (broadleaf), plantain (buckhorn), purslane, ragweed, sheepsorrel, shepherd's purse, speedwell, thistle

Rate	
lb/acre	lb/1000 sq ft
150-175	3.4 - 4

Application Directions

Application Directions

Apply ProScope fertilizer with LockUp™ using a drop or rotary-type spreader designed to apply granules. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly. Do not use spreaders that apply material in narrow concentrated bands. Avoid skips or overlaps as poor weed control or turfgrass injury may occur. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at right angles to the first.

To optimize product performance, mow the turfgrass two to three days before applying ProScope fertilizer with LockUp™. If the turfgrass is under moisture stress, a thorough sprinkler irrigation of at least 1/2 inch may be needed at least one to two days before application. For optimal results, apply ProScope fertilizer with LockUp™ in the morning when dew is present. If necessary, sprinkler irrigate with less than 1/4 inch of water just prior to application to maximize particle adherence to leaf surfaces. Avoid watering of treated area for 24 hours following application. Avoid making applications if rainfall is expected within 24 hours after treatment.

Application Techniques for Applying ProScope fertilizer with LockUp™

- When using a drop-type spreader, a splashboard mounted under the hopper will provide more even granule distribution.
- A chain fastened to the side of the spreader and allowed to drag on the soil surface can be used to mark the edge of the treated swath and help prevent skips or overlaps.
- For treating smaller areas, a hand held or push-type rotary applicator such as a whirlybird or cyclone unit is recommended. For hand held units, walk and turn the crank at a constant rate of speed.
- A shaker-type applicator made from a small container with holes punched in the bottom is recommended for small, difficult to treat areas. Carefully measure the amount of product needed to avoid over application.

Rates and Timing of Application

Apply 150 to 175 lb of ProScope fertilizer with LockUp™ per acre (3.4 to 4 lb per 1000 sq ft) per application to established turfgrass. Possible re-treatment in approximately 30 days may be necessary if turfgrass is heavily infested with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed 350 lb of ProScope fertilizer with LockUp™ per acre (8 lb per 1000 sq ft) per year because damage to turfgrass may occur. Do not make more than two applications per season.

Note: When applying to warm season turfgrass under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration. It is recommended to reduce application rates by 50% during hot, dry conditions to avoid possible dehydration or browning of turfgrass.

Best weed control results are obtained when ProScope fertilizer with LockUp™ is applied to small, actively growing weeds in the spring or early fall. Extreme growing conditions such as drought or near freezing temperatures prior to, at, or following time of application may reduce weed control. Broadleaf weed species germinate at different times.

Terms and Conditions of Use

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitation of Remedies.

Warranty Disclaimer

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. TO THE EXTENT PERMITTED BY LAW, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by buyer.

Limitation of Remedies

To the extent permitted by law, the exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

(1) Refund of purchase price paid by buyer or user for product bought, or (2) Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. To the extent permitted by law, in no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of the Seller or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.regulatory-info-lebsea.com>